

# Some descriptive set theory related to the Lebesgue density theorem

A. Andretta<sup>1</sup>   R. Camerlo<sup>2</sup>

<sup>1</sup>Dipartimento di Matematica  
Università di Torino

<sup>2</sup>Dipartimento di Matematica  
Politecnico di Torino

Wien 2009/6/23

# The category algebra.

Work in some perfect Polish space, e.g.  $\omega^2$ .

$\mathcal{B}$  is the collection of all sets with the property of Baire,

$\text{MGR}$  is the ideal of meager sets,

$$\mathcal{B}/\text{MGR} \cong \text{BOR}/\text{MGR} = \text{CAT}$$

$\text{CAT}$  is unique up-to isomorphism, i.e. it does not depend on the Polish space. The map

$$\rho: \text{CAT} \rightarrow \text{RO}$$

is a selector, and  $\text{CAT}$  can be identified with the collection of all regular open sets.

$\text{CAT}$  is a Polish space.

# The measure algebra.

$\mu$  a continuous probability Borel measure on some perfect Polish space, e.g. the usual Lebesgue measure on  $\omega^2$ .

MEAS is the collection of all sets measurable sets,

NULL is the ideal of measure-zero sets,

$$\text{MEAS}/\text{NULL} \cong \text{BOR}/\text{NULL} = \text{MALG}$$

MALG is unique up-to isomorphism, i.e. it does not depend on  $\mu$ .

MALG is a Polish space:

$$\delta([A], [B]) = \mu(A \triangle B)$$

## Definition

$x$  has density  $r \in [0; 1]$  in  $A$  if

$$d_A(x) \stackrel{\text{def}}{=} \lim_{n \rightarrow \infty} \frac{\mu(A \cap \mathbf{N}_{x \uparrow n})}{\mu(\mathbf{N}_{x \uparrow n})} = r.$$

## Theorem (Lebesgue)

Let  $A \subseteq {}^\omega 2$  be Lebesgue measurable. Then

$$\Phi(A) = \{x \in {}^\omega 2 \mid x \text{ has density 1 in } A\}$$

is Lebesgue measurable, and  $\mu(A \triangle \Phi(A)) = 0$ .

In other words:  $d_A$  agrees with  $\chi_A$  almost everywhere.

If  $\mu(A \triangle B) = 0$  then  $\Phi(A) = \Phi(B)$ , so

$$\Phi: \text{MALG} \rightarrow \text{MEAS}$$

is a selector. This is the analogue of  $\rho: \text{CAT} \rightarrow \text{RO}$ .

### Question

What is the complexity of  $\Phi(A)$ ?

## Definition

The localization of  $A$  at  $s$  is

$$A_{[s]} = \{x \in {}^\omega 2 \mid s \hat{\ } x \in A\}$$

Thus  $s \hat{\ } A_{[s]} = A \cap N_s$ .

## Trivial observation

$$\mu(A \triangle B) = 0 \Leftrightarrow \forall s \in {}^{<\omega} 2 (\mu(A_{[s]}) = \mu(B_{[s]}))$$

Thus a measure class  $[A]$  is completely determined by the map  $s \mapsto \mu(A_{[s]})$

Since

$$x \in \Phi(A) \Leftrightarrow \forall k \exists n \forall m \geq n \left( \mu(A_{[x \uparrow m]}) \geq 1 - 2^{-k-1} \right)$$

then

### Proposition (Folklore)

For all measurable  $A$

$$\Phi(A) \in \Pi_3^0.$$

### Question

Is  $\Pi_3^0$  optimal?

- $A \subseteq B \Rightarrow \Phi(A) \subseteq \Phi(B)$ ,
- $\Phi(A \cap B) = \Phi(A) \cap \Phi(B)$ ,
- $\bigcup_{i \in I} \Phi(A_i) \subseteq \Phi(\bigcup_{i \in I} A_i)$ ,
- if  $A$  is open, then  $A \subseteq \Phi(A)$  .

## Definition

$$\mathcal{T} = \{A \in \text{MEAS} \mid A \subseteq \Phi(A)\}$$

is the density topology. It is finer than the usual topology.



## Theorem (Scheinberg 1971, Oxtoby 1971?)

$A = \Phi(A)$  if and only if  $A$  is open and regular in  $\mathcal{T}$ .

$$\Phi: \text{MALG} \rightarrow \text{RO}_{\mathcal{T}}$$

- $\text{NULL} = \text{MGR}_{\mathcal{T}}$  (Oxtoby, 1971)
- $\mathcal{T}$  is neither first countable, nor second countable, nor Lindelöf, nor separable.
- $\mathcal{T}$  is Baire.

Recall that  $\Phi(A)$  is always  $\Pi_3^0$ .

### Theorem

There is a closed  $C$  such that  $\Phi(C)$  is complete  $\Pi_3^0$

Clearly

$$\text{Int}(A) \subseteq \Phi(A) \subseteq \text{Cl}(A).$$

and  $A = \Phi(A)$  if  $A$  is clopen.

### Question

Can  $\Phi(A)$  be something other than clopen or complete  $\Pi_3^0$ ?

Yes!

## Definition

$A$  is Wadge reducible to  $B$

$$A \leq_w B$$

just in case  $A = f^{-1}(B)$  for some continuous  $f: {}^\omega 2 \rightarrow {}^\omega 2$ .

$A \equiv_w B$  iff  $A \leq_w B \wedge B \leq_w A$ .

The equivalence classes  $[A]_w$  are called Wadge degrees.

## Theorem

$$\forall A \in \Pi_3^0 \exists C \in \Pi_1^0 (\Phi(C) \equiv_w A)$$

Recall that  $d_A(x) = 0, 1$  for *almost* all  $x$ .

### Definition

A set  $A$  is dualistic (or Manichæan) if  $d_A(x) = 0, 1$  for all  $x$ .  
 $\mathcal{M}$  is the Boolean algebra of all dualistic sets.

Clearly being dualistic depends on the equivalence class of  $A$ , so

$$A \in \mathcal{M} \Leftrightarrow \Phi(A) \in \mathcal{M}.$$

### Fact

$A = \Phi(A)$  is dualistic iff  $A$  is  $\mathcal{T}$ -clopen, i.e.,

$$\mathcal{M} \cap \text{ran}(\Phi) = \Delta_1^0\text{-}\mathcal{T}$$

## Proposition

$\forall A \in \text{MEAS} (A \in \mathcal{M} \Rightarrow \Phi(A) \in \Delta_2^0)$ .

## Theorem

$\forall A \in \Delta_2^0 \exists C \in \Pi_1^0 (\Phi(C) \equiv_W A \wedge \Phi(C) \in \mathcal{M})$ .

## Proposition

$\Phi$  is Borel (as a map from  $\text{MALG}$  to the set of codes for  $\Pi_3^0$  sets).

## Sketch of the proof for $\Pi_3^0$ completeness

- $T$  a pruned tree such that  $[T]$  has positive measure and empty interior. Thus  $\neg[T] = \bigcup_n \mathbf{N}_{t_n}$ .
- $n < m \Rightarrow \text{lh}(t_n) < \text{lh}(t_m)$  and  $\exists^\infty n (\text{lh}(t_n) + 1 < \text{lh}(t_{n+1}))$ .
- For all  $t \in T$  there is a shortest  $s \supset t$  such that  $s \notin T$ .  
 $s$  is the target of  $t$ .
- Let  $\tau(t) = \text{lh}(\text{target of } t) - \text{lh}(t)$ ,  $\tau: T \rightarrow \omega \setminus \{0\}$ .
- For  $x \in [T]$ ,

$$x \in \Phi([T]) \Leftrightarrow \lim_{n \rightarrow \infty} \tau(x \upharpoonright n) = \infty.$$

## Sketch of the proof for $\Pi_3^0$ completeness, ctd.

The set

$$P = \{z \in {}^{\omega \times \omega} 2 \mid \forall m \forall^\infty n z(n, m) = 0\}.$$

is complete  $\Pi_3^0$ .

Given  $a: n \times n \rightarrow 2$  construct a node  $\varphi(a) \in T$  so that

$$a \subset b \Rightarrow \varphi(a) \subset \varphi(b),$$

and

$${}^{\omega \times \omega} 2 \rightarrow [T], \quad z \mapsto \bigcup_n \varphi(z \upharpoonright n \times n)$$

witnesses  $P \leq_W \Phi([T])$ .

## Sketch of the proof for $\Pi_3^0$ completeness, ctd.

Let  $a: (n+1) \times (n+1) \rightarrow 2$ . (Say  $n = 4$ )

Case 1:

$a_{0,4}$	$a_{1,4}$	$a_{2,4}$	$a_{3,4}$	<b>0</b>
$a_{0,3}$	$a_{1,3}$	$a_{2,3}$	$a_{3,3}$	<b>0</b>
$a_{0,2}$	$a_{1,2}$	$a_{2,2}$	$a_{3,2}$	<b>0</b>
$a_{0,1}$	$a_{1,1}$	$a_{2,1}$	$a_{3,1}$	<b>0</b>
$a_{0,0}$	$a_{1,0}$	$a_{2,0}$	$a_{3,0}$	<b>0</b>

Then pick  $t \supset \varphi(a \upharpoonright n \times n)$  such that

$$\tau(t) \geq \max \{n+1, \tau(\varphi(a \upharpoonright n \times n))\}.$$



## Sketch of the proof for $\Pi_3^0$ completeness, ctd.

Let  $a: (n+1) \times (n+1) \rightarrow 2$ . (Say  $n = 4$ )

Case 2:


$a_{0,4}$	$a_{1,4}$	$a_{2,4}$	$a_{3,4}$	<b><math>a_{4,4}</math></b>
$a_{0,3}$	$a_{1,3}$	$a_{2,3}$	$a_{3,3}$	<b><math>a_{4,3}</math></b>
$a_{0,2}$	$a_{1,2}$	$a_{2,2}$	$a_{3,2}$	<b><math>a_{4,2}</math></b>
$a_{0,1}$	$a_{1,1}$	$a_{2,1}$	$a_{3,1}$	<b>0</b>
$a_{0,0}$	$a_{1,0}$	$a_{2,0}$	$a_{3,0}$	<b>0</b>

Then pick  $t \supset \varphi(a \upharpoonright n \times n)$  such that

$$\tau(t) = 3.$$


## The Wadge hierarchy on $\omega^2$ .

- A set  $A$  (or degree) is self dual if  $A \equiv_W \neg A$ . Otherwise it is non-self-dual.
- Self-dual and non-self-dual pairs alternate.
- At all limit levels there is a non-self-dual pair.



# How to construct larger degrees.

Given  $f: \omega \rightarrow \omega \setminus \{0\}$  and sets  $A_0, A_1, \dots$  consider the set




## How to construct larger degrees.

If  $\exists^\infty n (f(n) \geq 2)$  and the  $A_n$ s are  $\mathcal{T}$ -regular, i.e.  $\Phi(A_n) = A_n$  then so is  $\text{Rake}^-(f; (A_n)_n)$ . Moreover

- if  $A = A_0 = A_1 = \dots$  are self-dual, then  $\text{Rake}^-(f; (A_n)_n)$  is non-self-dual and immediately above  $A$ ,
- if  $A_0 <_W A_1 <_W A_2 <_W \dots$  then  $\text{Rake}^-(f; (A_n)_n)$  is non-self-dual and immediately above the  $A_n$ s.

Note that the rake  $\text{Rake}^-(f; (A_n)_n)$  has no pole, i.e.,  $0^{(\infty)}$  does not belong to this set. In order to construct the dual degrees we need another kind of rake, a pole and densely packed tines.

# How to construct larger degrees.



## How to construct larger degrees.

If  $\lim_n f(n) = \infty$  then and the  $A_n$ s are  $\mathcal{T}$ -regular, i.e.  $\Phi(A_n) = A_n$  then so is  $\text{Rake}^+(f; (A_n)_n)$ . Moreover

$$\text{Rake}^+(f; (A_n)_n) \equiv_W \neg \text{Rake}^-(f; (A_n)_n).$$

If  $A$  and  $B$  are  $\mathcal{T}$ -regular then so is

$$A \oplus B = 0 \hat{\wedge} A \cup 1 \hat{\wedge} B.$$

Arguing this way, we can climb up to  $\Delta_2^0$ .

## Jumping $\omega_1$ levels.


Wadge defined two operations  $A^{\natural}$  and  $A^{\flat}$  on subsets of the *Baire space*

$$A^{\natural} = \left\{ s_0^+ \smallfrown 0 \smallfrown s_1^+ \smallfrown 0 \smallfrown \dots \smallfrown s_n^+ \smallfrown 0 \smallfrown x^+ \mid n \in \omega, s_i \in {}^{<\omega}\omega, x \in A \right\}$$

$$A^{\flat} = A^{\natural} \cup \{x \in {}^{\omega}\omega \mid \exists^{\infty} n (x(n) = 0)\}$$

where  $s^+$  and  $x^+$  are the sequences obtained from  $s$  and  $x$  by adding a 1 to all entries.

The idea is that  $A^{\natural}$  is the union of  $\omega$  many layers of the form



# Jumping $\omega_1$ levels.

## Theorem (Wadge)

If  $A$  is self-dual, then  $A^{\flat}$  and  $A^{\flat\flat}$  form a non-self-dual pair and

$$\|A^{\flat}\|_{\mathbb{W}} = \|A^{\flat\flat}\|_{\mathbb{W}} = \|A\|_{\mathbb{W}} \cdot \omega_1.$$

The operations  $A^{\flat}$  and  $A^{\flat\flat}$  together with the (analogs of) the Rake operations, are sufficient to construct sets of rank  $< \omega_1^{\omega_1}$ , i.e. the  $\Delta_3^0$  sets.



## Jumping $\omega_1$ levels.

An analogue of  $A^+$ .

- $\overline{s \cap i} = \overline{s} \cap ii$ , for  $s \in {}^{<\omega}2$ .
- $\overline{x} = \bigcup_n \overline{x \upharpoonright n}$ , for  $x \in {}^\omega 2$ .
- Replace  $A$  with  $\{\overline{x} \mid x \in A\}$ , but...
- Does not work, since  $\{\overline{x} \mid x \in {}^\omega 2\}$  is of measure 0!
- The cure: enlarge  $\{\overline{x} \mid x \in A\}$  like  $\text{Rake}^-$  was enlarged to  $\text{Rake}^+$ . The resulting set is called  $\text{Plus}(A)$ .
- In fact we construct  $\text{Plus}(A; r)$  (with  $r \in (0; 1)$ ) so that  $\mu \left( \text{Plus}(A; r) \upharpoonright_{[\overline{s}]} \right) \geq r$  for all  $s$ .
- If  $A$  is  $\mathcal{T}$ -regular (i.e.,  $A = \Phi(A)$ ), then so is  $\text{Plus}(A; r)$ .


## Jumping $\omega_1$ levels.

Construct  $\text{Nat}(A)$  and  $\text{Flat}(A)$ : they are the analogs of  $A^{\natural}$  and  $A^{\flat}$ , and have rank  $\|A\|_{\mathbb{W}} \cdot \omega_1$ .

Using the operations  $\text{Nat}(A)$ ,  $\text{Flat}(A)$ ,  $\text{Rake}^- A$ ,  $\text{Rake}^+ A$ , and  $\oplus$  it is possible to construct a closed sets  $C$  such that  $\Phi(C)$  is of any given Wadge degree in  $\Delta_3^0$ .

# Nat( $A$ )

Fix  $0 < r < 1$ . Nat( $A$ ) is composed of  $\omega$ -many layers




- If  $x$  settles inside a layer, then  $x = s \hat{\ } \bar{y}$  and the density of  $x$  in Nat( $A$ ) will be 'similar' to the density of  $y$  in  $A$ .
- Every time we climb to a higher level, the density drops momentarily to  $\leq 1/2$ . So if  $x$  climbs infinitely many layers, then  $x$  will not have density 1 in Nat( $A$ ).

## Flat( $A$ )

Fix  $0 < r_0 < r_1 < r_2 < \dots \rightarrow 1$ .

Flat( $A$ ) is the set is composed of  $\omega$ -many layers



- If  $x$  settles inside a layer, then  $x = s^{\wedge} \bar{y}$  and the density of  $x$  in Flat( $A$ ) will be 'similar' to the density of  $y$  in  $A$ .
- In the layer  $n$ , the density will always be  $\geq r_n$ . So if  $x$  climbs infinitely many layers, then  $x$  will have density 1 in Flat( $A$ ).