

Topological dynamics of unordered Ramsey structures

by

Moritz Müller (Wien) and András Pongrác (Palaiseau)

Abstract. We investigate the connections between Ramsey properties of Fraïssé classes \mathcal{K} and the universal minimal flow $M(G_{\mathcal{K}})$ of the automorphism group $G_{\mathcal{K}}$ of their Fraïssé limits. As an extension of a result of Kechris, Pestov and Todorcevic (2005) we show that if the class \mathcal{K} has finite Ramsey degree for embeddings, then this degree equals the size of $M(G_{\mathcal{K}})$. We give a partial answer to a question of Angel, Kechris and Lyons (2014) showing that if \mathcal{K} is a relational Ramsey class and $G_{\mathcal{K}}$ is amenable, then $M(G_{\mathcal{K}})$ admits a unique invariant Borel probability measure that is concentrated on a unique generic orbit.

1. Introduction. With a Fraïssé class of finite structures \mathcal{K} one can associate in a natural way a topological group $G_{\mathcal{K}}$, namely, the automorphism group of the Fraïssé limit of \mathcal{K} . For example, the Fraïssé limit of finite-dimensional vector spaces over a fixed finite field F is the \aleph_0 -dimensional vector space $V_{\infty, F}$ over F with automorphism group $\mathrm{GL}(V_{\infty, F})$. The groups of the form $G_{\mathcal{K}}$ are precisely the Polish groups that are non-Archimedean in the sense that they have a basis at the identity consisting of open subgroups [2].

In [14] Kechris, Pestov and Todorcevic developed a “duality theory” [13, §4(A)] linking finite combinatorics of \mathcal{K} with topological dynamics of $G_{\mathcal{K}}$; more precisely, it links combinatorial properties of \mathcal{K} with properties of the universal minimal $G_{\mathcal{K}}$ -flow $M(G_{\mathcal{K}})$. For groups of the form $G_{\mathcal{K}}$ the flow $M(G_{\mathcal{K}})$ is an inverse limit of metrisable $G_{\mathcal{K}}$ -flows (cf. [14, T1.5]), and in many interesting cases it is metrisable itself ⁽¹⁾. If so, $M(G_{\mathcal{K}})$ either has the size of the continuum or else is finite [14, §1(E)]. An extreme case is when $M(G_{\mathcal{K}})$ is a single point, that is, $G_{\mathcal{K}}$ is extremely amenable. It is shown in [14] that for ordered \mathcal{K} this happens if and only if \mathcal{K} is Ramsey. For

2010 *Mathematics Subject Classification*: Primary 05C55; Secondary 37B05, 03C15.

Key words and phrases: finite flow, Ramsey, amenable, measure concentration.

⁽¹⁾ Very recently, Zucker [25, T8.7] showed that $M(G_{\mathcal{K}})$ is metrisable if and only if every structure in \mathcal{K} has finite Ramsey degree in \mathcal{K} .

example, $V_{\infty, F}$ together with the so-called “canonical order” has an extremely amenable automorphism group.

We give a characterisation of $M(G_{\mathcal{K}})$ having an arbitrary finite cardinality in terms of Ramsey properties of \mathcal{K} . Namely, we use Ramsey degrees [8, 9, 10] and show that $M(G_{\mathcal{K}})$ has finite size d if and only if \mathcal{K} has Ramsey degree d (Theorem 3.1). We do not assume \mathcal{K} to be ordered, but use Ramsey degrees *for embeddings* instead (see e.g. [18, 3]). These coincide with the usual Ramsey degrees on rigid structures, so our characterisation generalises the above mentioned result of [14] and so does its proof.

Given an appropriate (unordered) class \mathcal{K} one can first produce a so-called reasonable order expansion \mathcal{K}^* whose Fraïssé limit expands the limit of \mathcal{K} by a (linear) order $<^*$. The group $G_{\mathcal{K}}$ acts naturally on orders and one gets a $G_{\mathcal{K}}$ -flow $X_{\mathcal{K}^*}$ as the orbit closure $\overline{G_{\mathcal{K}} \cdot <^*}$. Again, as shown in [14], minimality of this flow corresponds to a combinatorial property of \mathcal{K}^* called the ordering property (cf. [18]), and indeed $X_{\mathcal{K}^*}$ is $M(G_{\mathcal{K}})$ if and only if \mathcal{K}^* is additionally Ramsey ⁽²⁾. Moreover, the Ramsey degree of $A \in \mathcal{K}$ equals the number of non-isomorphic order expansions it has in \mathcal{K}^* ([14, §10], [19, §4]).

For example, the universal minimal $\text{GL}(V_{\infty, F})$ -flow is the orbit closure of the canonical order. This canonical order is forgetful in the sense that any finite-dimensional F -vector space gets, up to isomorphism, only one order expansion, so Ramsey degrees are 1 in this case. The Ramsey degrees *for embeddings*, on the other hand, are unbounded. In general, the relationship between the two degrees is not trivial. We show that if a Ramsey class in a relational language has finite Ramsey degree for embeddings, then this degree must be a power of 2 (Theorem 3.11).

Recently, Angel, Kechris and Lyons [1] extended the duality theory to other important properties of $M(G_{\mathcal{K}})$, namely whether or not there is a (unique) $G_{\mathcal{K}}$ -invariant Borel probability measure on $M(G_{\mathcal{K}})$. In this case, the group $G_{\mathcal{K}}$ is called amenable (uniquely ergodic), and this happens if and only if all minimal $G_{\mathcal{K}}$ -flows admit such a (unique) measure [1, P8.1]. For example, $\text{GL}(V_{\infty, F})$ is uniquely ergodic.

The $G_{\mathcal{K}}$ -flows $X_{\mathcal{K}^*}$ have a generic (i.e. comeager) orbit $G_{\mathcal{K}} \cdot <^*$ which is in fact dense G_{δ} [1, P14.3]. In many examples, a $G_{\mathcal{K}}$ -invariant measure on $M(G_{\mathcal{K}})$, if it exists, turns out to be concentrated on this generic orbit. However, answering a question in [1, Q15.3], Zucker [24, T1.2] showed that the measure on $M(\text{GL}(V_{\infty, F}))$ is *not* concentrated on the generic orbit.

We show that such counterexamples rely on the language containing function symbols. More precisely, we show that if \mathcal{K} is Ramsey over a relational language and $G_{\mathcal{K}}$ is amenable, then $G_{\mathcal{K}}$ is uniquely ergodic and the unique

⁽²⁾ See [20] for a discussion of how to characterise universality alone.

$G_{\mathcal{K}}$ -invariant Borel probability measure on $M(G_{\mathcal{K}})$ is indeed concentrated on a dense G_{δ} orbit (Theorem 4.1).

2. Preliminaries

2.1. Notation. For $k \in \mathbb{N}$ we let $[k]$ denote $\{0, \dots, k-1\}$, and understand $[0] = \emptyset$. If X, Y are sets, f a function from X to Y , $n \in \mathbb{N}$ and $Z \subseteq X^n$, we write $f(Z)$ for the set $\{f(\bar{x}) \mid \bar{x} \in Z\}$ where $f(\bar{x})$ denotes the tuple $(f(x_0), \dots, f(x_{n-1}))$ for $\bar{x} = (x_0, \dots, x_{n-1}) \in X^n$. For $X_0 \subseteq X$ we let $f|X_0$ denote the restriction of f to X_0 ; for a relation Z as above, $Z|X_0$ denotes $Z \cap (X_0^n)$. The identity on X is denoted by id_X .

2.2. Fraïssé theory. Fix a countable language L . We let A, B, \dots range over (L -)structures. The distinction between structures and their universes is blurred notationally. We speak of *relational* structures and classes of structures if the underlying language L is relational. We write $A \leq B$ to indicate that there exists an embedding from A into B , and we let B^A denote the set of embeddings from A into B .

The *age* $\text{Age}(F)$ of a structure F is the class of finitely generated structures which embed into F . A structure F is *locally finite* if its finitely generated substructures are finite. For $A \in \text{Age}(F)$ we call F *A -homogeneous* if for all $a, a' \in F^A$ there is $g \in \text{Aut}(F)$ such that $g \circ a = a'$. If F is A -homogeneous for all $A \in \text{Age}(F)$, it is (*ultra*-)homogeneous.

A structure F is *Fraïssé* if it is countably infinite, locally finite and homogeneous. The age $\mathcal{K} := \text{Age}(F)$ of a Fraïssé structure F :

- is *hereditary*: for all A, B , if $A \leq B$ and $B \in \mathcal{K}$, then $A \in \mathcal{K}$;
- has *joint embedding*: for all $A, B \in \mathcal{K}$ there is $C \in \mathcal{K}$ such that both $A \leq C$ and $B \leq C$;
- has *amalgamation*: for all $A, B_0, B_1 \in \mathcal{K}$ and $a_0 \in B_0^A, a_1 \in B_1^A$ there are $C \in \mathcal{K}$ and $b_0 \in C^{B_0}, b_1 \in C^{B_1}$ such that $b_0 \circ a_0 = b_1 \circ a_1$.

A class \mathcal{K} of finite structures that has these three properties, contains countably many structures up to isomorphism, and for every $n \in \mathbb{N}$ contains a structure (with universe) of size at least n , is a *Fraïssé class*. The following is well-known [22, T4.4.4]:

THEOREM 2.1 (Fraïssé, 1954). *For every Fraïssé class \mathcal{K} there exists a Fraïssé structure F with age \mathcal{K} .*

A standard back-and-forth argument shows that the structure F in Theorem 2.1 is unique up to isomorphism; it is called the *Fraïssé limit* of \mathcal{K} and denoted by $\text{Flim}(\mathcal{K})$.

We mention some standard examples:

EXAMPLES 2.2. The Fraïssé limit of the class of linear orderings is the rational order $(\mathbb{Q}, <)$. The Fraïssé limit of the class of finite Boolean algebras is the countable atomless Boolean algebra B_∞ . The Fraïssé limit of the class of finite graphs is the random graph R . The Fraïssé limit of the class of finite vector spaces over a fixed finite field F is the vector space $V_{\infty, F}$ of dimension \aleph_0 over F .

We refer to [6, 7, 16] for surveys on homogeneous structures.

2.3. Ramsey degrees. Write $\binom{B}{A}$ for the set of substructures of B which are isomorphic to A . Note that $\binom{B'}{A} \subseteq \binom{C}{A}$ whenever $B' \in \binom{C}{B}$. If $k, d \in \mathbb{N}$ then $C \rightarrow (B)_{k,d}^A$ means that for every colouring $\chi : \binom{C}{A} \rightarrow [k]$ there exists $B' \in \binom{C}{B}$ such that $|\chi(\binom{B'}{A})| \leq d$. The *Ramsey degree* of A in a class \mathcal{K} of structures is the least $d \in \mathbb{N}$ such that for all $B \in \mathcal{K}$ and $k \geq 2$ there is $C \in \mathcal{K}$ such that $C \rightarrow (B)_{k,d}^A$, provided that such a d exists; otherwise it is ∞ . Taking the supremum over $A \in \mathcal{K}$ gives the Ramsey degree of \mathcal{K} , and the Ramsey degree of a structure F is understood to be the Ramsey degree of $\text{Age}(F)$; if this degree is 1, then \mathcal{K} resp. F are simply called *Ramsey*.

EXAMPLES 2.3. $(\mathbb{Q}, <)$, B_∞ and $V_{\infty, F}$ are Ramsey [14]. The random graph R has Ramsey degree ∞ ; indeed, a finite graph G has Ramsey degree $|G|!/|\text{Aut}(G)|$ in the class of finite graphs [14, §10].

We refer to the surveys [11, 17] for more information on Ramsey theory.

2.4. Topological dynamics. With a Fraïssé class \mathcal{K} we associate the topological group

$$G_{\mathcal{K}} := \text{Aut}(\text{Flim}(\mathcal{K})),$$

the identity having basic neighbourhoods

$$G_{(A)} := \{g \in G_{\mathcal{K}} \mid g \upharpoonright A = \text{id}_A\}$$

for all finite substructures A of $\text{Flim}(\mathcal{K})$. For any topological group G a G -flow is a continuous action $a : G \times X \rightarrow X$ of G on a compact Hausdorff space X . When the action is understood we shall refer to X as a G -flow and write $g \cdot x$ or gx for $a(g, x)$. For $Y \subseteq X$ we write $G \cdot Y := \bigcup_{g \in G} gY = \bigcup_{y \in Y} Gy$ where $Gy := \{gy \mid g \in G\}$ denotes the *orbit* of y and $gY := \{gy \mid y \in Y\}$.

EXAMPLE 2.4. Let $G = \text{Aut}(F)$ for a countable structure F . The *space of linear orders* (on F) is $LO := \{R \subseteq F^2 \mid R \text{ is a linear order on } F\}$ with topology given by basic open sets $\{R \mid R_0 \subseteq R\}$ for R_0 a linear order on a finite subset A of F . This space is compact and Hausdorff, and is a G -flow with respect to $(g, R) \mapsto g(R)$, the *logic action* of G on LO .

A subset $Y \subseteq X$ is G -invariant if $G \cdot Y \subseteq Y$. Closed G -invariant subsets Y are G -flows with respect to the restriction of the action. Such G -flows are *subflows* of X . The flow X is *minimal* if X and \emptyset are its only subflows, that

is, if and only if every orbit is dense. By Zorn's lemma, every G -flow contains a minimal subflow. A *homomorphism* (resp. *isomorphism*) of a G -flow X into another Y is a continuous (resp. continuous bijective) G -map $\pi : X \rightarrow Y$; being a G -map means that $\pi(g \cdot x) = g \cdot \pi(x)$ for all $g \in G, x \in X$.

The following is well-known (cf. [23, §3]).

THEOREM 2.5. *For every Hausdorff topological group G there exists a minimal G -flow $M(G)$ which is universal in the sense that for every minimal G -flow Y there is a homomorphism from X into Y . Any two universal minimal G -flows are isomorphic.*

An interesting case is when $|M(G)| = 1$, or equivalently, every G -flow X has a fixed point, i.e. an $x \in X$ such that $G \cdot x = \{x\}$. In this case G is called *extremely amenable*. Being *amenable* means that there exists a (Borel probability) measure μ on $M(G)$ which is G -invariant (i.e. $\mu(X) = \mu(g \cdot X)$ for every Borel $X \subset M(G)$ and $g \in G$). If there is exactly one such measure then G is *uniquely ergodic*. It is shown in [1, P8.1] that for a uniquely ergodic G in fact every minimal G -flow has a unique G -invariant measure.

We refer to [14, §1] for a survey on universal minimal flows.

2.5. Duality theory. Let $<$ be a binary relation symbol. A class \mathcal{K}^* of finite $L \cup \{<\}$ -structures is *ordered* if each of its members has the form $(A, <^A)$ for a (linear) order $<^A$ (on A) and some finite L -structure A ; the order $<^A$ is called a \mathcal{K}^* -admissible one (cf. [18]).

The following is [14, T4.8].

THEOREM 2.6. *Assume that \mathcal{K}^* is an ordered Fraïssé class. Then $G_{\mathcal{K}^*}$ is extremely amenable if and only if \mathcal{K}^* is Ramsey.*

Let $\mathcal{K} := \{A \mid (A, <^A) \in \mathcal{K}^*\}$ be the L -reduct of \mathcal{K}^* ; \mathcal{K}^* is *reasonable* if for all $A, B \in \mathcal{K}$, all $a \in B^A$ and all \mathcal{K}^* -admissible orders $<^A$ on A there is a \mathcal{K}^* -admissible order $<^B$ on B such that $a(<^A) \subseteq <^B$, i.e. $a \in (B, <^B)^{(A, <^A)}$.

LEMMA 2.7. *Let \mathcal{K} be a Fraïssé class and let $F = \text{Flim}(\mathcal{K})$. Then $\mathcal{K}^* = \text{Age}(F, R)$ is reasonable for every order R on F .*

Proof. Let $A, B \in \mathcal{K}$, $a \in B^A$ and $<^A$ be a \mathcal{K}^* -admissible order on A . Let $a_0 \in (F, R)^{(A, <^A)}$ and $b \in F^B$. In particular, $a_0 \in F^A$ and $b \circ a \in F^A$, and then by homogeneity of F there exists an $\alpha \in \text{Aut}(F)$ such that $\alpha \circ b \circ a = a_0$. We define

$$<^B := (b^{-1} \circ \alpha^{-1})(R \upharpoonright (\alpha \circ b)(B)).$$

We need to show that $a^{-1}(<^B \upharpoonright a(A)) = <^A$. We have

$$\begin{aligned} a^{-1}(<^B \upharpoonright a(A)) &= a^{-1}((b^{-1} \circ \alpha^{-1})(R \upharpoonright (\alpha \circ b)(B)) \upharpoonright a(A)) \\ &= a^{-1}((b^{-1} \circ \alpha^{-1})(R \upharpoonright (\alpha \circ b)(a(A)))) = a_0^{-1}(R \upharpoonright a_0(A)) = <^A. \end{aligned}$$

The last equality holds as $a_0 \in (F, R)^{(A, <^A)}$. ■

The following is [14, P5.2, T10.8]. Recall that LO denotes the space of orders (Example 2.4).

THEOREM 2.8. *Let \mathcal{K}^* be a reasonable ordered Fraïssé class in the language $L \cup \{<\}$, and \mathcal{K} its L -reduct.*

- (1) *The class \mathcal{K} is Fraïssé and $\text{Flim}(\mathcal{K}^*) = (\text{Flim}(\mathcal{K}), <^*)$ for some linear order $<^*$.*
- (2) *Let $X_{\mathcal{K}^*} := \overline{G_{\mathcal{K}} \cdot <^*}$ be the orbit closure of $<^*$ in the logic action of $G_{\mathcal{K}}$ on LO . Then $X_{\mathcal{K}^*}$ is the universal minimal $G_{\mathcal{K}}$ -flow if and only if \mathcal{K}^* is Ramsey and has the ordering property.*

That \mathcal{K}^* has the *ordering property* means that for all $A \in \mathcal{K}$ there is a $B \in \mathcal{K}$ such that $(A, <^A) \leq (B, <^B)$ for all \mathcal{K}^* -admissible orders $<^A$ on A and $<^B$ on B .

In [1] Kechris et al. showed that a certain quantitative version of the ordering property characterises unique ergodicity for so-called Hrushovski classes. Here, we shall only need the following [1, P9.2].

PROPOSITION 2.9. *Let \mathcal{K}^* be a reasonable ordered Fraïssé class which is Ramsey and satisfies the ordering property, and let \mathcal{K} be its L -reduct. Then $G_{\mathcal{K}}$ is amenable (resp. uniquely ergodic) if and only if there exists a consistent random \mathcal{K}^* -admissible ordering $(R_A)_{A \in \mathcal{K}}$ (resp. and for any other consistent random \mathcal{K}^* -admissible ordering $(R'_A)_{A \in \mathcal{K}}$, R_A and R'_A have the same distribution for every $A \in \mathcal{K}$).*

Indeed, if $(R_A)_{A \in \mathcal{K}}$ is a consistent random \mathcal{K}^ -admissible ordering, then there is a $G_{\mathcal{K}}$ -invariant Borel probability measure μ on $X_{\mathcal{K}^*}$ such that for every $A \in \mathcal{K}$ and every \mathcal{K}^* -admissible ordering $<$ on A we have ⁽³⁾ $\mu(U(<)) = \text{Pr}[R_A = <]$ where*

$$U(<) := \{R \in X_{\mathcal{K}^*} \mid R \upharpoonright A = <\}.$$

A *random \mathcal{K}^* -admissible ordering* is a family $(R_A)_{A \in \mathcal{K}}$ of random variables such that each R_A takes values in the set of \mathcal{K}^* -admissible orders on A . It is *consistent* if for all $A, B \in \mathcal{K}$ and $a \in B^A$ the random variables $a^{-1}(R_B \upharpoonright \text{im}(a))$ and R_A have the same distribution.

EXAMPLES 2.10. In [14, §6] the reader can find constructions of reasonable ordered Fraïssé classes \mathcal{K}^* whose reduct \mathcal{K} is any of the classes mentioned in Examples 2.2; in all these cases \mathcal{K}^* is Ramsey and has the ordering property. By Theorem 2.6 one sees that the automorphism groups of $(\mathbb{Q}, <)$ and of certain ordered versions of B_∞ , R , $V_{\infty, F}$ are extremely amenable [14]. Theorem 2.8 allows us to calculate the universal minimal flows of the auto-

⁽³⁾ Given a random variable we always use Pr to denote the probability measure of its underlying probability space.

morphism groups of B_∞ , R and $V_{\infty,F}$. The group $\text{Aut}(B_\infty)$ is not amenable, while $\text{Aut}(R)$ and $\text{Aut}(V_{\infty,F})$ are uniquely ergodic [1].

3. Automorphism groups with finite universal minimal flows.

Theorem 2.6 characterises the condition that the universal minimal flow has size 1. In this section we provide a similar characterisation for the condition that it has an arbitrary finite size. To this end we consider Ramsey degrees for embeddings. The main result in this section reads:

THEOREM 3.1. *Let $d \in \mathbb{N}$ and \mathcal{K} be a Fraïssé class. The following are equivalent:*

- (1) $M(G_{\mathcal{K}})$ has size at most d ;
- (2) the Ramsey degree for embeddings of \mathcal{K} is at most d .

We start with some preliminary observations concerning finite universal minimal flows in Section 3.1. In Section 3.2 we define Ramsey degrees for embeddings and discuss their relationship to Ramsey degrees. The results proved in Sections 3.1 and 3.2 are mainly folklore. In Section 3.3 we prove the result above and in Section 3.4 we note some corollaries.

3.1. Finite universal minimal flows

LEMMA 3.2. *Let G be a topological Hausdorff group and $d \in \mathbb{N}$. Then $M(G)$ has size at most d if and only if every non-empty G -flow has an orbit of size at most d .*

Proof. Assume that $|M(G)| \leq d$, and let X be a non-empty G -flow. Then there is a minimal subflow X' of X and a homomorphism π of $M(G)$ onto X' . Thus $|X'| \leq d$.

Conversely, if every non-empty G -flow has an orbit of size at most d , then so does $M(G)$. Since $M(G)$ is minimal, this orbit is dense in $M(G)$, so it is equal to $M(G)$ by finiteness. ■

LEMMA 3.3. *Let G be a topological Hausdorff group and H an extremely amenable closed subgroup of G with finite index. Then H is a normal clopen subgroup of G and $M(G)$ is isomorphic to the action of G on G/H by left multiplication.*

Proof. Clearly, a closed subgroup of finite index is open. We first show that G/H is the universal minimal G -flow. Since H is open, G/H is discrete, and as $|G : H|$ is finite, G/H is compact. Hence, G/H is a G -flow. It is minimal, because G acts transitively on G/H . If Y is an arbitrary G -flow, then its restriction to H is an H -flow, so it has a fixed point $y \in Y$. Then $gH \mapsto gy$ is a homomorphism from G/H into Y .

As gHg^{-1} is a closed subgroup of finite index for every $g \in G$, so is $H' = H \cap gHg^{-1}$. As above, we see that G/H' is a minimal G -flow. By

universality of G/H there exists a surjection from G/H onto G/H' , so $|G : H'| \leq |G : H|$. Thus $H = gHg^{-1}$ for every $g \in G$, that is, H is normal. ■

This lemma gives the forward direction of the following proposition. The proof of the converse has been explained to the authors by Todor Tsankov. It adapts an argument due to Kechris and Sokić [15, Appendix 3].

PROPOSITION 3.4. *Let G be a topological Hausdorff group and $d \in \mathbb{N}$. Then $M(G)$ has size d if and only if G has an extremely amenable, open, normal subgroup of index d .*

Proof. The backward direction follows from Lemma 3.3. Conversely, assume that $X := M(G)$ has size d . For $x \in X$ let $H_x \leq G$ be the stabiliser of x . Then there is a bijection between the set of left cosets of H_x and the orbit $G \cdot x$. Since $G \cdot x$ is finite, we have $G \cdot x = \overline{G \cdot x}$, so $G \cdot x = X$ by minimality. Hence, $|G : H_x| = |X| = d$. As H_x is closed and of finite index, so is $N := \bigcap_{x \in X} H_x$, and hence N is clopen. Since N is the pointwise stabiliser of X , it is normal. Let Y be a minimal N -flow. Let $\tau : G/N \rightarrow G$ be a function with $\tau(hN) \in hN$. Define $a : G \times G/N \rightarrow N$ by setting

$$a(g, hN) := \tau(hN)^{-1} \cdot g^{-1} \cdot \tau(ghN).$$

A straightforward calculation shows that a satisfies the so-called cocycle identity, that is, for all $g_1, g_2, h \in G$,

$$(1) \quad a(g_1g_2, hN) = a(g_2, hN) \cdot a(g_1, g_2hN).$$

We can construct an action of G on $(G/N \times Y)$ by

$$(g, (hN, y)) \mapsto (ghN, a(g, hN)^{-1} \cdot y).$$

That this indeed defines a group action follows directly from (1). The action is continuous and $(G/N \times Y)$ is compact, so $(G/N \times Y)$ is a G -flow.

Let $h \in G$ and $y \in Y$ be arbitrary. We show that

$$(2) \quad Y(h, y) := \{a(n, hN)^{-1} \cdot y \mid n \in N\} \text{ is dense in } Y.$$

Indeed, as N is normal, we have $Y(h, y) = \tau(hN)^{-1} \cdot N \cdot \tau(hN) \cdot y = N \cdot y$. Since Y is a minimal N -flow, the orbit $N \cdot y$ is dense in Y .

The orbit $G \cdot (hN, y)$ contains $N \cdot (ghN, y')$ for every $g \in G$ and $y' := a(g, hN)^{-1} \cdot y$. But $N \cdot (ghN, y') = \{(nghN, a(n, hN)^{-1} \cdot y') \mid n \in N\} = \{ghN\} \times Y(h, y')$, where the last equality holds because N is normal. So the orbit $G \cdot (hN, y)$ contains $\bigcup_{g \in G} (\{gN\} \times Y(h, y_g))$ for certain y_g 's, and this set is dense in $(G/N \times Y)$ by (2). Thus $(G/N \times Y)$ is a minimal G -flow.

By the universality of X there exists a surjection from X onto $(G/N \times Y)$. In particular, $|G/N \times Y| \leq d$. By definition of N we have $|G : N| \geq d$, so $|Y| = 1$ and $|G/N| = d$. This means that N is extremely amenable and has index d in G . ■

EXAMPLE 3.5. For $d \in \mathbb{N}$ let G^* be the automorphism group of $(\mathbb{Q}, <, 0, \dots, d-1)$, the structure with universe \mathbb{Q} that interprets for all $i \in [d]$ a constant by i and a binary relation symbol $<$ by the rational order. Let G be the group generated by G^* and the permutation $\alpha = (0 \ 1 \ \dots \ d-1)$. This is a closed subgroup of the group of all permutations of \mathbb{Q} , so $G = G_{\mathcal{K}}$ for some Fraïssé class \mathcal{K} (see e.g. [2]). Since α commutes with G , G^* is normal in G . Moreover, G^* has index d in G , and it follows from [5, L13] (see also [4, P24]) that G^* is extremely amenable. By Lemma 3.3, $|M(G)| = |G/G^*| = d$.

EXAMPLE 3.6. Let G be the automorphism group of $(\mathbb{Q}, E_d, <)$ where $<$ is the rational order and E_d is an equivalence relation with d classes each of which is dense in $(\mathbb{Q}, <)$. Let H be the subgroup of G consisting of those automorphisms that preserve each of these classes. It is shown in [14, T8.4] that H is extremely amenable and of index $d!$ in G . By Lemma 3.3, $|M(G)| = |G/H| = d!$.

3.2. Ramsey degrees for embeddings. Let $k, d \in \mathbb{N}$ and let \mathcal{K} be a class of finite structures. Then $C \hookrightarrow (B)_{k,d}^A$ means that for every colouring $\chi : C^A \rightarrow [k]$ there exists a $b \in C^B$ such that $|\chi(b \circ B^A)| \leq d$. Naturally, here $b \circ B^A$ denotes $\{b \circ a \mid a \in B^A\}$. The *Ramsey degree for embeddings* of A in \mathcal{K} is the least $d \in \mathbb{N}$ such that for all $B \in \mathcal{K}$ and $k \geq 2$ there is a $C \in \mathcal{K}$ such that $C \hookrightarrow (B)_{k,d}^A$, provided that such a d exists; otherwise it is ∞ . Taking the supremum over $A \in \mathcal{K}$ gives the Ramsey degree for embeddings of \mathcal{K} . If this degree is 1, we call \mathcal{K} *Ramsey for embeddings*.

LEMMA 3.7. *Let $d \in \mathbb{N}$, \mathcal{K} be a Fraïssé class, $F = \text{Flim}(\mathcal{K})$ and $A \in \mathcal{K}$. The Ramsey degree for embeddings of A in \mathcal{K} is at most d if and only if $F \hookrightarrow (B)_{k,d}^A$ for all $B \in \mathcal{K}$ and $k \geq 2$.*

Proof. Assume that the Ramsey degree for embeddings of A in \mathcal{K} is at most d . Let $B \in \mathcal{K}$, $k \geq 2$ and $\chi : F^A \rightarrow [k]$. We are looking for $b' \in F^B$ such that $|\chi(b' \circ B^A)| \leq d$. Choose $C \in \mathcal{K}$ such that $C \hookrightarrow (B)_{k,d}^A$. Choose a $c \in F^C$ and let $\chi' : C^A \rightarrow [k]$ map $a \in C^A$ to $\chi(c \circ a)$. By $C \hookrightarrow (B)_{k,d}^A$ there is a $b \in C^B$ such that $|\chi'(b \circ B^A)| \leq d$, i.e. $|\chi(c \circ b \circ B^A)| \leq d$. Then $b' := c \circ b \in F^B$ is as desired.

Assume that there is an $A \in \mathcal{K}$ whose Ramsey degree for embeddings is bigger than d . Choose $B \in \mathcal{K}$ and $k \geq 2$ such that for every finite substructure C of F there is a colouring $\chi : C^A \rightarrow [k]$ which is *good for C* , i.e. $|\chi(b \circ B^A)| > d$ for all $b \in C^B$. The set $G(C) := \{\chi \in [k]^{F^A} \mid \chi \upharpoonright C^A \text{ is good for } C\}$ is non-empty and closed in $[k]^{F^A}$ carrying the product topology with $[k]$ being discrete. Given finitely many such sets $G(C_1), \dots, G(C_n)$ their intersection contains the non-empty set $G(C)$ where C is the substructure generated by $C_1 \cup \dots \cup C_n$ in F (note that C is finite by local finiteness

of F). Since $[k]^{F^A}$ is compact, $\bigcap_C G(C) \neq \emptyset$ where C ranges over the finite substructures of F . Any $\chi \in \bigcap_C G(C)$ is good for F , so $F \not\rightarrow (B)_{k,d}^A$. ■

We shall need the following result of Nešetřil [18, T3.2]. We include the short proof.

LEMMA 3.8. *Let \mathcal{K} be a hereditary class of finite structures with joint embedding. If \mathcal{K} is Ramsey for embeddings, then it has amalgamation.*

Proof. Let $A, B_0, B_1 \in \mathcal{K}$ and $a_0 \in B_0^A, a_1 \in B_1^A$. Let $B \in \mathcal{K}$ and $b_0 \in B^{B_0}, b_1 \in B^{B_1}$. Choose $C \in \mathcal{K}$ with $C \hookrightarrow (B)_{4,1}^A$. We claim that there exist $e_0 \in C^{B_0}$ and $e_1 \in C^{B_1}$ such that $e_0 \circ a_0 = e_1 \circ a_1$. Consider the following colouring $\chi : C^A \rightarrow P(\{0, 1\})$: for $a \in C^A$ the colour $\chi(a) \subseteq \{0, 1\}$ contains $i \in \{0, 1\}$ if and only if there exists an $e \in C^{B_i}$ such that $e \circ a_i = a$. Choose $b \in C^B$ such that $\chi(b \circ B^A)$ contains precisely one colour. Then this colour is $\{0, 1\}$, because for $i \in \{0, 1\}$ we have $i \in \chi(b \circ b_i \circ a_i)$ and $b \circ b_i \circ a_i \in b \circ B^A$. Let $a \in B^A$. Then $\chi(b \circ a) = \{0, 1\}$, thus there are $e_0 \in C^{B_0}$ and $e_1 \in C^{B_1}$ such that $e_0 \circ a_0 = a = e_1 \circ a_1$. ■

REMARK 3.9. Clearly, $C \hookrightarrow (B)_{k,d}^A$ is equivalent to $C \rightarrow (B)_{k,d}^A$ when A is rigid (i.e. $\text{Aut}(A) = \{\text{id}_A\}$). In particular, the Ramsey degree and the Ramsey degree for embeddings coincide for rigid structures. The following proposition generalises this observation (see [25, P4.4]).

PROPOSITION 3.10. *Let $d \in \mathbb{N}$, and let \mathcal{K} be a class of finite structures. Let $A \in \mathcal{K}$ and $\ell = |\text{Aut}(A)|$. The Ramsey degree for embeddings of A in \mathcal{K} is $d \cdot \ell$ if and only if the Ramsey degree of A in \mathcal{K} is d .*

In Section 4.3 we shall prove the following.

THEOREM 3.11. *Let \mathcal{K} be a relational Fraïssé class which is Ramsey. Then the Ramsey degree for embeddings of \mathcal{K} is infinite or a finite power of 2.*

We refer to Examples 4.5 for some natural examples of relational Fraïssé classes which are Ramsey and have infinite Ramsey degree for embeddings.

3.3. Proof of Theorem 3.1. The theorem is a consequence of the following two propositions which in fact establish something stronger.

We say that a class of finite structures \mathcal{D} is *cofinal* in another such class \mathcal{K} if for all $A \in \mathcal{K}$ there exists $B \in \mathcal{D}$ such that $A \leq B$.

PROPOSITION 3.12. *Let $d \in \mathbb{N}$ and let \mathcal{K} be a Fraïssé class. Assume that the class of structures with Ramsey degree for embeddings at most d in \mathcal{K} is cofinal in \mathcal{K} . Then $M(G_{\mathcal{K}})$ has size at most d .*

Proof. Write $G := G_{\mathcal{K}}$ and $F := \text{Flim}(\mathcal{K})$. Let $A \in \mathcal{K}$, $a_0 \in F^A$ and write $A_0 := \text{im}(a_0)$. Consider the map $\Phi : G \rightarrow F^A$, $g \mapsto g \circ a_0$. By homogeneity

of F , Φ is surjective. For all $g, h \in G$ we have

$$g \circ a_0 = h \circ a_0 \Leftrightarrow gG_{(A_0)} = hG_{(A_0)}.$$

Hence, Φ induces a bijection e from $G/G_{(A_0)}$ onto F^A . Observe that

$$(3) \quad g \circ e(hG_{(A_0)}) = g \circ (h \circ a_0) = (gh) \circ a_0 = e((gh)G_{(A_0)}).$$

CLAIM 1. *Assume that A has Ramsey degree for embeddings at most d in \mathcal{K} . Let $k \in \mathbb{N}$ and let $f : G \rightarrow [k]$ be constant on each $gG_{(A_0)} \subseteq G$ for $g \in G$. Then for every finite $H \subseteq G$ there exists $g \in G$ such that $|f(gH)| \leq d$.*

Proof. The function f induces a function \tilde{f} from $G/G_{(A_0)}$ to $[k]$. Note that $\tilde{f} \circ e^{-1} : F^A \rightarrow [k]$. There is a finite substructure $B \subseteq F$ such that

$$(4) \quad \{e(hG_{(A_0)}) \mid h \in H\} \subseteq B^A.$$

By Lemma 3.7 there is a $b \in F^B$ such that $|(\tilde{f} \circ e^{-1})(b \circ B^A)| \leq d$. By homogeneity of F there is a $g \in G$ such that $g \circ \text{id}_B = b$. We show that g is as desired, namely $f(gh) \in (\tilde{f} \circ e^{-1})(b \circ B^A)$ for every $h \in H$:

$$f(gh) = \tilde{f}((gh)G_{(A_0)}) = \tilde{f} \circ e^{-1}(e((gh)G_{(A_0)})) = \tilde{f} \circ e^{-1}(g \circ e(hG_{(A_0)}))$$

where the last equality follows from (3). By (4), $g \circ e(hG_{(A_0)}) \in g \circ B^A = b \circ B^A$, and our claim follows. \dashv

For $n \in \mathbb{N}$, $n \geq 1$, consider \mathbb{R}^n with the Euclidean norm $\|\cdot\|$. For $\varepsilon > 0$ and $x \in \mathbb{R}^n$ let

$$B_\varepsilon(x) := \{y \in \mathbb{R}^n \mid \|x - y\| < \varepsilon\}.$$

As a topological group G carries its *left* uniformity, that is, the uniformity with basic entourages $\{(g, h) \mid g^{-1}h \in G_{(A)}\}$ for $A \in \text{Age}(M)$, $A \subseteq M$.

CLAIM 2. *Let n be a positive integer, $f : G \rightarrow \mathbb{R}^n$ be left uniformly continuous and bounded, $H \subseteq G$ be finite and ε be a positive real. Then there are $g \in G$ and $h_0, \dots, h_{d-1} \in H$ such that*

$$(5) \quad f(gH) \subseteq \bigcup_{\nu < d} B_\varepsilon(f(gh_\nu)).$$

Proof. By left uniform continuity of f there is a finite substructure $A' \subseteq F$ such that $\|f(g) - f(g')\| < \varepsilon/6$ for all $g, g' \in G$ with $gG_{(A')} = g'G_{(A')}$. By our cofinality assumption, there exist $A'' \in \mathcal{K}$ and $a' \in (A'')^{A'}$ such that A'' has Ramsey degree for embeddings at most d in \mathcal{K} . Since F is homogeneous, there is an embedding $a'' \in F^{A''}$ such that $a'' \circ a' = \text{id}_{A'}$. Hence, the image A of a'' has Ramsey degree for embeddings at most d in \mathcal{K} , and $A' \subseteq A \subseteq F$. Thus $G_{(A)} \subseteq G_{(A')}$, so for all $g, g' \in G$ with $gG_{(A)} = g'G_{(A)}$,

$$(6) \quad \|f(g) - f(g')\| < \varepsilon/6.$$

We claim that there exists a function $\tilde{f} : G \rightarrow \mathbb{R}^n$ such that

- (a) $\text{im}(\tilde{f})$ is finite;
- (b) \tilde{f} is constant on $gG_{(A)}$ for every $g \in G$;
- (c) $\|f(g) - \tilde{f}(g)\| < \varepsilon/2$ for every $g \in G$.

By (a) and (b) we can apply Claim 1 and obtain some $g \in G$ such that $|\tilde{f}(gH)| \leq d$. Choose $h_0, \dots, h_{d-1} \in H$ such that $\tilde{f}(gH) = \{\tilde{f}(gh_\nu) \mid \nu < d\}$. To verify (5), let $h \in H$ be given. We have to show that there exists $\nu < d$ such that $\|f(gh) - f(gh_\nu)\| < \varepsilon$. Indeed, this holds for $\nu < d$ such that $\tilde{f}(gh) = \tilde{f}(gh_\nu)$, because by (c) both $\|f(gh) - \tilde{f}(gh_\nu)\| = \|f(gh) - \tilde{f}(gh)\| < \varepsilon/2$ and $\|f(gh_\nu) - \tilde{f}(gh_\nu)\| < \varepsilon/2$.

Thus, it remains to find \tilde{f} with properties (a)–(c).

As f is bounded, its image is contained in a compact subset of \mathbb{R}^n . Choose finitely many points $y_\nu \in \mathbb{R}^n$, $\nu < k'$, such that this compact set is covered by $\bigcup_{\nu < k'} B_{\varepsilon/6}(y_\nu)$. Assume that precisely the first $k \leq k'$ balls $B_{\varepsilon/6}(y_\nu)$ contain a point from the image of f . For $\nu < k$ choose $\hat{\nu} \in G$ such that $f(\hat{\nu}) \in B_{\varepsilon/6}(y_\nu)$. Then $\bigcup_{\nu < k} B_{\varepsilon/3}(f(\hat{\nu}))$ covers the image of f . Hence, for every $g \in G$ we can choose $\nu_g < k$ such that

$$(7) \quad \|f(g) - f(\hat{\nu}_g)\| < \varepsilon/3.$$

Let $c : G \rightarrow G$ be a selector for the partition $\{gG_{(A)} \mid g \in G\}$ of G , that is, for all $g, g' \in G$ we have $c(g) \in gG_{(A)}$, and $c(g) = c(g')$ if and only if $gG_{(A)} = g'G_{(A)}$. Define

$$\tilde{f}(g) := f(\widehat{\nu_{c(g)}}).$$

Then \tilde{f} satisfies (a) and (b). For all $g \in G$ we have $c(g) \in gG_{(A)}$, so $gG_{(A)} = c(g)G_{(A)}$ and thus $\|f(g) - f(c(g))\| < \varepsilon/6$ by (6). As $\|f(c(g)) - f(\widehat{\nu_{c(g)}})\| < \varepsilon/3$ by (7), we conclude that \tilde{f} satisfies (c). \dashv

We aim to show that every G -flow has an orbit of size at most d (Lemma 3.2). So let X be a G -flow. We are looking for some $x_0 \in X$ such that

$$(8) \quad |G \cdot x_0| \leq d.$$

Recall that the compact Hausdorff space X carries a unique uniformity compatible with its topology. Suppose f is a uniformly continuous function from X into \mathbb{R}^n for some $n \geq 1$. For each $x \in X$ define $f_x : G \rightarrow \mathbb{R}^n$ by

$$f_x(g) := f(g^{-1} \cdot x).$$

Then f_x is left uniformly continuous. This follows from the well-known fact that for every $x \in X$ the map $g \mapsto g^{-1} \cdot x$ is left uniformly continuous (see e.g. [21, L2.1.5]).

With a triple (H, f, ε) for a finite subset $H \subseteq G$, and a bounded, uniformly continuous function $f : X \rightarrow \mathbb{R}^n$, and a real $\varepsilon > 0$ we associate the

set

$$Y(H, f, \varepsilon) := \left\{ x \in X \mid \exists h_0, \dots, h_{d-1} \in H : f_x(H) \subseteq \bigcup_{\nu < d} \overline{B_\varepsilon(f_x(h_\nu))} \right\}.$$

Since H is finite, $Y(H, f, \varepsilon)$ is a finite union of closed sets of the form $\{x \in X \mid f_x(H) \subseteq C\}$ for $C \subseteq \mathbb{R}^n$ closed, and consequently $Y(H, f, \varepsilon)$ is closed.

CLAIM 3. *The family of closed sets $Y(H, f, \varepsilon)$ with H, f, ε as above has the finite intersection property.*

Proof. For $j < \ell$ let $H_j \subseteq G$ be finite, $\varepsilon_j > 0$ and $f^j : X \rightarrow \mathbb{R}^{n_j}$ for $n_j \geq 1$. Set $H := \bigcup_{j < \ell} H_j$, $\varepsilon := \min_{j < \ell} \varepsilon_j$, $n := \sum_{j < \ell} n_j$ and define $f : X \rightarrow \mathbb{R}^n$ by $f(x) := f^0(x) * \dots * f^{\ell-1}(x)$ where $*$ denotes concatenation. Then f is uniformly continuous and bounded.

Let $x \in X$ be arbitrary. Since $f_x : G \rightarrow \mathbb{R}^n$ is left uniformly continuous, Claim 2 applies, and there exist $g \in G$ and $h_0, \dots, h_{d-1} \in H$ such that $f_x(gH) \subseteq \bigcup_{\nu < d} B_\varepsilon(f_x(gh_\nu))$. In other words,

$$(9) \quad \forall h \in H \exists \nu < d : f(h^{-1}g^{-1}x) \in B_\varepsilon(f(h_\nu^{-1}g^{-1}x)).$$

Any $y \in \mathbb{R}^n$ can be written as $y[0] * \dots * y[\ell-1]$, where $y[j] \in \mathbb{R}^{n_j}$ for all $j < \ell$. In this notation, $f_x(g)[j] = f_x^j(g)$ for all $g \in G$, $x \in X$, $j < \ell$. Clearly, $f_x(g) \in B_\varepsilon(y)$ implies $f_x(g)[j] \in B_\varepsilon(y[j])$ for all $y \in \mathbb{R}^n$, $j < \ell$. Writing $x_0 := g^{-1}x$, we see that (9) yields

$$\forall j < \ell \forall h \in H_j \exists \nu < d : f(h^{-1}g^{-1}x)[j] = f_{x_0}^j(h) \in B_\varepsilon(f_{x_0}^j(h_\nu)).$$

Since $\varepsilon \leq \varepsilon_j$, we obtain

$$\forall j < \ell : f_{x_0}^j(H_j) \subseteq \bigcup_{\nu < d} B_{\varepsilon_j}(f_{x_0}^j(h_\nu)).$$

Thus, $x_0 \in \bigcap_{j < \ell} Y(H_j, f^j, \varepsilon_j) \neq \emptyset$. \dashv

By Claim 3 and since X is compact, there exists an x_0 in the intersection of all the sets $Y(H, f, \varepsilon)$ with (H, f, ε) a triple as above. We claim that x_0 satisfies (8). Assume otherwise that there are $g_0, \dots, g_d \in G$ such that g_0x_0, \dots, g_dx_0 are pairwise distinct. Choose $f : X \rightarrow [0, 1] \subseteq \mathbb{R}^1$ uniformly continuous such that $f(g_\nu x_0) = \nu/d$ for all $\nu \leq d$. Then $x_0 \notin Y(\{g_\nu^{-1} \mid \nu \leq d\}, f, \varepsilon)$ for a small enough $\varepsilon > 0$, a contradiction. \blacksquare

PROPOSITION 3.13. *Let $d \in \mathbb{N}$, let F be countable and locally finite, $G := \text{Aut}(F)$, and $A \in \text{Age}(F)$ such that F is A -homogeneous. If $M(G)$ has size at most d , then $F \hookrightarrow (B)_{k,d}^A$ for all $B \in \text{Age}(F)$ and $k \geq 2$.*

Proof. Assume that $|M(G)| \leq d$, $B \in \text{Age}(F)$, $k \geq 2$ and let $\chi_0 : F^A \rightarrow [k]$ be a colouring. Note that $[k]^{F^A}$ is compact Hausdorff in the product topology with $[k]$ being discrete. The group G acts continuously on $[k]^{F^A}$ by the shift $(g, \chi) \mapsto g \cdot \chi$, where $g \cdot \chi$ colours $a \in F^A$ by $\chi(g^{-1} \circ a)$. Consider

the orbit closure $\overline{G \cdot \chi_0}$ of χ_0 . By Lemma 3.2, the induced action of G on $\overline{G \cdot \chi_0}$ has an orbit of size at most d , that is, there exist $\chi_1 \in \overline{G \cdot \chi_0}$ and $\psi_0, \dots, \psi_{d-1} \in \overline{G \cdot \chi_0}$ such that $G \cdot \chi_1 = \{\psi_i \mid i < d\}$.

Let $b \in F^B$. Observe that $b \circ B^A$ is a finite subset of F^A . Since $\chi_1 \in \overline{G \cdot \chi_0}$, there exists a $g \in G$ such that $g \cdot \chi_0$ and χ_1 agree on $b \circ B^A$. Note that $g^{-1} \circ b \in F^B$, so it remains to show that $|\chi_0(g^{-1} \circ b \circ B^A)| \leq d$. We fix some $a_0 \in F^A$, and claim that for all $a \in g^{-1} \circ b \circ B^A$ there exists a $\nu < d$ such that $\chi_0(a) = \psi_\nu(a_0)$. To see this, let $a \in g^{-1} \circ b \circ B^A \subseteq F^A$ and choose $h \in G$ so that $h \circ a_0 = a$. Such an h exists since F is A -homogeneous. Then

$$\chi_0(a) = (g \cdot \chi_0)(g \circ a) = \chi_1(g \circ a) = \chi_1((gh) \circ a_0) = ((gh)^{-1} \cdot \chi_1)(a_0),$$

where the second equality follows from $g \circ a \in b \circ B^A$ and the choice of g . As $(gh)^{-1} \cdot \chi_1 \in G \cdot \chi_1$, and by the choice of χ_1 , there exists $\nu < d$ such that $(gh)^{-1} \cdot \chi_1 = \psi_\nu$. Thus $\chi_0(a) = \psi_\nu(a_0)$ as claimed. ■

Proof of Theorem 3.1. (1) \Rightarrow (2). Write $F = \text{Flim}(\mathcal{K})$, and let $A \in \mathcal{K} = \text{Age}(F)$. Then F and A satisfy the assumptions of Proposition 3.13, and so $F \hookrightarrow (B)_{k,d}^A$ for all $B \in \mathcal{K}$ and $k \geq 2$. Now apply Lemma 3.7.

(2) \Rightarrow (1). By Proposition 3.12. ■

3.4. Corollaries

COROLLARY 3.14. *Let $d \in \mathbb{N}$ and \mathcal{K} be a Fraïssé class. The following are equivalent ⁽⁴⁾:*

- (1) *The class of structures with Ramsey degree for embeddings at most d in \mathcal{K} is cofinal in \mathcal{K} .*
- (2) *\mathcal{K} has Ramsey degree for embeddings at most d .*

Proof. Assume (1). By Proposition 3.12 we have $|M(G_{\mathcal{K}})| \leq d$. As $F := \text{Flim}(\mathcal{K})$ is Fraïssé, Proposition 3.13 implies $F \hookrightarrow (B)_{k,d}^A$ for all $A, B \in \mathcal{K}$. Then Lemma 3.7 implies (2). ■

It is noted in [14, §1(D)] that a separable metrisable group G is extremely amenable, i.e. $M(G)$ has size 1, if and only if every metrisable G -flow has a fixed point. In this context it might be of interest to note:

COROLLARY 3.15. *Let $d \in \mathbb{N}$ and \mathcal{K} be a Fraïssé class. The following are equivalent:*

- (1) *$M(G_{\mathcal{K}})$ has size at most d .*
- (2) *Every continuous action of $G_{\mathcal{K}}$ on the Cantor space has an orbit of size at most d .*

Proof. (1) implies (2) by Lemma 3.2. Conversely, assume (2). Let $A \in \mathcal{K}$ and write $F := \text{Flim}(\mathcal{K})$. Then F and A satisfy the assumptions of Propo-

⁽⁴⁾ Meanwhile, Zucker gave a simpler, direct proof of this corollary [25, P4.7].

sition 3.13. In the proof of this proposition we only require the following for $G_{\mathcal{K}}$: for all $k \geq 2$ and all $\chi_0 \in [k]^{F^A}$, the shift action of $G_{\mathcal{K}}$ restricted to $\overline{G_{\mathcal{K}} \cdot \chi_0}$ has a small orbit. But $[k]^{F^A}$ is homeomorphic to the Cantor space, and the restricted shift is a continuous action on this space. Thus (2) suffices to carry out this proof, and we conclude that $F \hookrightarrow (B)_{k,d}^A$ for all $B \in \mathcal{K} = \text{Age}(F)$. By Lemma 3.7 every $A \in \mathcal{K}$ has Ramsey degree for embeddings at most d in \mathcal{K} . Then Proposition 3.12 implies (1). ■

4. Measure concentration. We say that a probability measure is *concentrated* on any set of measure 1. In this section we prove the following.

THEOREM 4.1. *Let \mathcal{K} be a relational Fraïssé class which is Ramsey. If $G_{\mathcal{K}}$ is amenable, then it is uniquely ergodic and the (unique) $G_{\mathcal{K}}$ -invariant Borel probability measure on $M(G_{\mathcal{K}})$ is concentrated on a (unique) dense $G_{\mathcal{S}}$ orbit.*

In Section 4.1 we construct a forgetful order expansion using the Ramsey property, in Section 4.2 we prove Theorem 4.1, and the final Section 4.3 contains some observations concerning the ω -categorical case and a proof of (a stronger version of) Theorem 3.11.

4.1. Forgetful order expansions. An ordered class \mathcal{K}^* of finite structures in the language $L \cup \{<\}$ is called *forgetful* if for all $A, B \in \mathcal{K}$ and \mathcal{K}^* -admissible orderings $<^A, <^B$ on A, B respectively, we have $(A, <^A) \cong (B, <^B)$ whenever $A \cong B$; here \mathcal{K} denotes the L -reduct of \mathcal{K}^* .

For instance, the orderings of B_{∞} and $V_{\infty, F}$ mentioned in Examples 2.10 have forgetful ages (see [14, §6] for details). The following is easy to see (cf. [14, P5.6]).

LEMMA 4.2. *Let \mathcal{K}^* be a forgetful ordered class of finite structures in the language $L \cup \{<\}$ and \mathcal{K} its L -reduct. Then \mathcal{K}^* has the ordering property, and \mathcal{K}^* is Ramsey if and only if so is \mathcal{K} .*

Before showing that the Ramsey property ensures the existence of reasonable forgetful expansions, we present a well-known technical lemma. Informally, this lemma guarantees a monochromatic copy of a given B when copies of several different A_i are coloured simultaneously.

LEMMA 4.3. *Let \mathcal{K} be a Ramsey class. Let $n \in \mathbb{N}$, $k_0, \dots, k_{n-1} \in \mathbb{N}$, $A_0, \dots, A_{n-1}, B \in \mathcal{K}$. Then there exists a $C \in \mathcal{K}$ with the following property: for any family of colourings $\chi_i : \binom{C}{A_i} \rightarrow [k_i]$, $i \in [n]$, there exists a $B' \in \binom{C}{B}$ such that $\chi_i \upharpoonright B'$ is constant for all $i \in [n]$.*

Proof. Let $C_0 := B$, and for every $0 < i \leq n$ choose $C_i \in \mathcal{K}$ such that $C_i \rightarrow (C_{i-1})_{k_{i-1}, 1}^{A_{i-1}}$. Let $C := C_n$. Then by using the definition of the C_i and a straightforward induction on $j \in [n]$ we deduce that there is a

$C'_{n-1-j} \in \binom{C}{C_{n-1-j}}$ such that $\chi_i \upharpoonright C'_{n-1-j}$ is constant for all $i \in [n] \setminus [n-1-j]$. Setting $j = n-1$ yields B' as in the statement. ■

LEMMA 4.4. *Let \mathcal{K} be a Fraïssé class in the language L . If \mathcal{K} is Ramsey, then there exists a reasonable, forgetful ordered Fraïssé class \mathcal{K}^* in the language $L \cup \{<\}$ with L -reduct \mathcal{K} .*

Proof. Let $F := \text{Flim}(\mathcal{K})$ and consider the space LO of linear orders on F (cf. Example 2.4). Let (A, B) range over pairs with $A \in \mathcal{K}$ and $B \subseteq F$. Call $R \in LO$ order forgetful for (A, B) if $(A', R \upharpoonright A') \cong (A'', R \upharpoonright A'')$ for all $A', A'' \in \binom{B}{A}$.

CLAIM. *If $n \geq 1$ and $(A_0, B_0), \dots, (A_{n-1}, B_{n-1})$ are pairs as above with all $B_i \subseteq F$ finite, then there exists $R \in LO$ that is order forgetful for every (A_i, B_i) , $i \in [n]$.*

Proof. Choose $B \subseteq F$ finite such that $\bigcup_{i \in [n]} B_i \subseteq B$. It suffices to find an order which is order forgetful for every (A_i, B) , $i \in [n]$. Fix an arbitrary order $R \in LO$. For $i \in [n]$ let χ_i colour each $A'_i \in \binom{F}{A_i}$ by the isomorphism type of $(A'_i, R \upharpoonright A'_i)$, and let $k_i \in \mathbb{N}$ be the number of colours of χ_i . By Lemma 4.3 and homogeneity of F there exist $B' \subseteq F$ and $g \in \text{Aut}(F)$ such that B' is monochromatic and $g(B') = B$. By definition of the χ_i this means that R is order forgetful for (A_i, B') for all $i \in [n]$. Hence, $g(R)$ is order forgetful for all (A_i, B) , $i \in [n]$. ◻

For every $A \in \mathcal{K}$ and $B \subseteq F$ finite, the set of orders that are order forgetful for (A, B) is closed in LO . By the Claim and compactness, there exists $R \in LO$ which is order forgetful for all pairs (A, B) such that $A \in \mathcal{K}$ and $B \subseteq F$ is finite. Then R is order forgetful for (A, F) for every $A \in \mathcal{K}$. Equivalently, $\mathcal{K}^* := \text{Age}(F, R)$ is forgetful. To see that \mathcal{K}^* is Fraïssé, observe that \mathcal{K}^* is hereditary and has joint embedding. As \mathcal{K}^* is Ramsey by Lemma 4.2, it has amalgamation by Lemma 3.8 (and Remark 3.9; note that \mathcal{K}^* is rigid because it is ordered). According to Lemma 2.7, \mathcal{K}^* is reasonable. ■

EXAMPLES 4.5. The structures $F_1 := (\mathbb{Q}, \text{Betw})$, $F_2 := (\mathbb{Q}, \text{Cycl})$, $F_3 := (\mathbb{Q}, \text{Sep})$ and $F_4 := (\mathbb{Q}, =)$ are Ramsey (see [12] for definitions). If $<$ is the rational order, then $\mathcal{K}_i^* := \text{Age}((F_i, <))$ is forgetful with reduct $\mathcal{K}_i := \text{Age}(F_i)$. By Lemmas 2.7, 4.2 and Theorem 2.8, $M(G_{\mathcal{K}_i})$ is $\overline{G_{\mathcal{K}_i}} \cdot <$. Then $M(G_{\mathcal{K}_1})$ is the 2-element discrete space. Hence, by Theorem 3.1, \mathcal{K}_1 has Ramsey degree for embeddings 2. Theorem 2.8 also allows us to explicitly describe $M(G_{\mathcal{K}_i})$ for $i = 2, 3, 4$, and these have the size of the continuum. Hence, $\mathcal{K}_2, \mathcal{K}_3, \mathcal{K}_4$ have infinite Ramsey degree for embeddings.

EXAMPLES 4.6. Let \mathcal{K} be a Fraïssé class of digraphs such that there is a directed cycle in \mathcal{K} . Then there does not exist a forgetful ordered Fraïssé class with L -reduct \mathcal{K} : by forgetfulness, every directed edge in any $A \in \mathcal{K}$

would be ordered in the same way, and then a directed cycle contradicts transitivity of the order. For example, this applies to the age of the universal homogeneous digraph, the random tournament and the local order (see [16]).

4.2. Proof of Theorem 4.1. Let $F := \text{Flim}(\mathcal{K})$ and let L denote the relational language of \mathcal{K} . Since \mathcal{K} is assumed to be Ramsey, Lemma 4.4 applies and there is a reasonable forgetful ordered Fraïssé class \mathcal{K}^* in the language $L \cup \{<\}$ with L -reduct \mathcal{K} . By Lemma 4.2 and Theorem 2.8, $\text{Flim}(\mathcal{K}^*) = (F, <^*)$ for some order $<^*$, and $X_{\mathcal{K}^*} = \overline{G_{\mathcal{K}} \cdot <^*}$ is the universal minimal flow of $G_{\mathcal{K}}$.

Assume that $G_{\mathcal{K}}$ is amenable. In order to verify that $G_{\mathcal{K}}$ is uniquely ergodic, it suffices by Proposition 2.9 to show that for every consistent random ordering $(R_A)_{A \in \mathcal{K}}$ each random variable R_A is uniformly distributed. By forgetfulness, for any two \mathcal{K}^* -admissible orderings $<, <'$ on A there is an $\alpha \in \text{Aut}(A)$ such that $\alpha(<) = <'$, and then $\Pr[R_A = <'] = \Pr[\alpha^{-1} \circ R_A = <] = \Pr[R_A = <]$ where the latter equality follows from $(R_A)_A$ being consistent.

Let μ denote the unique $G_{\mathcal{K}}$ -invariant Borel probability measure on $X_{\mathcal{K}^*}$. Recall the notation $U(<)$ from Proposition 2.9. By this result, $U(<)$ and $U(<')$ have the same μ -measure whenever $<$ and $<'$ are \mathcal{K}^* -admissible orderings of the same finite subset of F .

An order $R \in X_{\mathcal{K}^*}$ is outside $G_{\mathcal{K}} \cdot <^*$ if and only if $(F, R) \not\cong (F, <^*)$, if and only if (F, R) is not homogeneous (cf. Section 2.2), if and only if there exist a finite $A \subseteq F$, some $(B, <^B) \in \mathcal{K}^*$ and $a \in (B, <^B)^{(A, <^* \upharpoonright A)}$ such that R is bad for $(B, <^B, a)$, meaning that there is no $b \in (F, R)^{(B, <^B)}$ with $b \circ a = \text{id}_A$. As the language of F is relational, we may assume that $B = \text{im}(a) \cup \{p\}$ with $p \in F \setminus \text{im}(a)$.

Observe that the set of orders $R \in X_{\mathcal{K}^*}$ which are bad for $(B, <^B, a)$ is closed. Hence, $X_{\mathcal{K}^*} \setminus G_{\mathcal{K}} \cdot <^*$ is F_{σ} , so $G_{\mathcal{K}} \cdot <^*$ is a dense G_{δ} orbit in $X_{\mathcal{K}^*}$ (see also [1, 14.3]). Since $X_{\mathcal{K}^*}$ is a Baire space, $G_{\mathcal{K}} \cdot <^*$ is clearly unique with this property. We prove that $\mu(G_{\mathcal{K}} \cdot <^*) = 1$. It suffices to show that for each $(B, <^B, a)$ with $B = \text{im}(a) \dot{\cup} \{p\}$ as above, the set $\mathcal{B} := \{R \in X_{\mathcal{K}^*} \mid R \text{ is bad for } (B, <^B, a)\}$ has μ -measure 0.

We construct a sequence $(\mathcal{U}_n)_{n \in \mathbb{N}}$ such that for all $n \in \mathbb{N}$:

- (a) \mathcal{U}_n is a cover of \mathcal{B} , i.e. $\mathcal{B} \subseteq \bigcup \mathcal{U}_n$;
- (b) every $U \in \mathcal{U}_n$ equals some $U(<')$ such that $<' \supseteq <^* \upharpoonright A$ is a \mathcal{K}^* -admissible order with $|\text{dom}(<')| = |A| + n$;
- (c) $\mu\left(\bigcup \mathcal{U}_{n+1}\right) \leq \frac{|A| + n}{|A| + n + 1} \cdot \mu\left(\bigcup \mathcal{U}_n\right)$.

Here, $\text{dom}(<')$ is the set linearly ordered by $<'$; note that (b) implies that $\text{dom}(<') \supseteq A$.

This finishes the proof: by (a) and (c) we have, for all $n \in \mathbb{N}$,

$$\mu(\mathcal{B}) \leq \mu\left(\bigcup \mathcal{U}_n\right) \leq \prod_{m < n} \frac{|A| + m}{|A| + m + 1} \cdot \mu\left(\bigcup \mathcal{U}_0\right) = \mu\left(\bigcup \mathcal{U}_0\right) \cdot \frac{|A|}{|A| + n} \xrightarrow{n} 0.$$

Set $\mathcal{U}_0 := \{U(<^* \upharpoonright A)\}$, and assume that \mathcal{U}_n is already defined. It suffices to find for every $U(<') \in \mathcal{U}_n$ some $p' \notin \text{dom}(<')$ and a family $(<_i)_{i \in I}$ such that

- (a') $\bigcup_{i \in I} U(<_i) \cap \mathcal{B} = U(<') \cap \mathcal{B}$;
- (b') for every $i \in I$, $<_i \supseteq <'$ is a \mathcal{K}^* -admissible order with $\text{dom}(<_i) = \text{dom}(<') \cup \{p'\}$;
- (c') $\mu\left(\bigcup_{i \in I} U(<_i)\right) \leq \frac{|A| + n}{|A| + n + 1} \cdot \mu(U(<'))$.

Write $A' := \text{dom}(<')$, and choose $R \in \mathcal{B} \cap U(<')$. Since $R \in \overline{G_{\mathcal{K}} \cdot <^*}$, there is a $g \in G_{\mathcal{K}}$ such that

$$(10) \quad g(<^*) \upharpoonright A' = R \upharpoonright A' = <'.$$

In particular, $g(<^*) \upharpoonright A = <' \upharpoonright A = <^* \upharpoonright A$ and $(A, <^* \upharpoonright A)$ is a substructure of $(F, g(<^*))$. Since $(F, g(<^*))$ is isomorphic to $(F, <^*)$, it is homogeneous, so there exists an embedding $b \in (F, g(<^*))^{(B, <^B)}$ with $b \circ a = \text{id}_A$. We set $p' := b(p)$ and claim that $p' \notin A'$. Otherwise, $\text{im}(b) \subseteq A'$, so $b \in (F, R)^{(B, <^B)}$ by (10), and this contradicts R being bad for $(B, <^B, a)$.

Let $<_0, \dots, <_{s-1}$ list the \mathcal{K}^* -admissible orders on $A' \cup \{p'\}$ extending $<'$, and note that $s \leq |A'| + 1$. Let $I \subseteq [s]$ consist of those $i < s$ such that $U(<_i) \cap \mathcal{B} \neq \emptyset$. Then (a') and (b') follow, and it remains to verify (c'). The sets $U(<_i)$, $i < s$, partition $U(<')$ and, as already noted, have pairwise equal μ -probability, so $\mu(U(<_i)) = \mu(U(<'))/s$. Thus

$$(11) \quad \mu\left(\bigcup_{i \in I} U(<_i)\right) = |I|/s \cdot \mu(U(<')).$$

There exists $i_0 < s$ with $<_{i_0} = g(<^*) \upharpoonright (A' \cup \{p'\})$. Since $b \in (F, g(<^*))^{(B, <^B)}$ has $\text{im}(b) \subseteq A' \cup \{p'\}$, we have $b \in (F, S)^{(B, <^B)}$ for every $S \in U(<_{i_0})$. Hence, no such S is bad for $(B, <^B, a)$, that is, $U(<_{i_0}) \cap \mathcal{B} = \emptyset$, so $i_0 \notin I$. Thus $|I| < s$. Since $|A'| = |A| + n$, we have $s \leq |A| + n + 1$, so $|I|/s \leq (s-1)/s \leq (|A| + n)/(|A| + n + 1)$. Hence, (c') follows from (11). This finishes the proof of Theorem 4.1.

REMARK 4.7. In the statement of Theorem 4.1 the condition that the language is relational cannot be omitted. According to Zucker's result [24, T1.2], the generic orbit of $M(\text{GL}(V_{\infty, F}))$ has measure 0. Unlike in the proof of Theorem 4.1, if the language of F contains function symbols, then it is not possible to extend the domain of the substructure A by just one point at a time. This algebraic nature of the structure would make it impossible

for us to have sufficient control over the number of admissible orders on a substructure generated by n elements and the set A . In particular, a sequence $(\mathcal{U}_n)_{n \in \mathbb{N}}$ as in the above proof cannot be constructed for the vector space. It would be interesting to know whether Theorem 4.1 holds for relational Fraïssé classes \mathcal{K} where every structure in \mathcal{K} has finite Ramsey degree in \mathcal{K} .

4.3. The ω -categorical case. Of particular interest are Fraïssé classes \mathcal{K} which have an ω -categorical Fraïssé limit $F := \text{Flim}(\mathcal{K})$. By the theorem of Ryll-Nardzewski (see e.g. [22, T4.3.1]) this happens e.g. if the language L of \mathcal{K} is finite and relational (cf. [22, T4.4.7]), and is equivalent to $G_{\mathcal{K}}$ being *oligomorphic*: for every $n \in \mathbb{N}$, $G_{\mathcal{K}}$ has only finitely many n -orbits. An n -orbit of $G_{\mathcal{K}}$ is an orbit of the *diagonal action* of $G_{\mathcal{K}}$ on F^n given by $g \cdot \bar{a} = g \cdot (a_0, \dots, a_{n-1}) := g(\bar{a}) = (g(a_0), \dots, g(a_{n-1}))$.

LEMMA 4.8. *Let \mathcal{K}^* be a reasonable ordered Fraïssé class in the language $L \cup \{<\}$ with L -reduct \mathcal{K} . Then $G_{\mathcal{K}^*}$ is oligomorphic if and only if so is $G_{\mathcal{K}}$.*

Proof. Let $F = \text{Flim}(\mathcal{K})$. By Theorem 2.8 we have $\text{Flim}(\mathcal{K}^*) = (F, <^*)$ for some order $<^*$ on F . As $G_{\mathcal{K}^*}$ is a subgroup of $G_{\mathcal{K}}$, it suffices to show that every orbit $T \subseteq F^n$ of $G_{\mathcal{K}}$ that consists of tuples with all different entries is the union of finitely many n -orbits of $G_{\mathcal{K}^*}$. Let $\bar{s} = (s_1, \dots, s_n)$ and $\bar{t} = (t_1, \dots, t_n)$ be tuples in T such that the unique extension of the partial isomorphism $s_1 \mapsto t_1, \dots, s_n \mapsto t_n$ to the substructures in F generated by \bar{s} and \bar{t} is a partial isomorphism of F^* . Then by homogeneity of F^* we find that \bar{s} and \bar{t} are in the same n -orbit of $G_{\mathcal{K}^*}$. As there are finitely many ways to define a (\mathcal{K}^* -admissible) order on the structure generated by a tuple in T , the claim follows. ■

LEMMA 4.9. *Let \mathcal{K}^* be a reasonable ordered Fraïssé class in the language $L \cup \{<\}$ with L -reduct \mathcal{K} . Assume that $G_{\mathcal{K}^*}$ is oligomorphic. If $G_{\mathcal{K}^*}$ is normal in $G_{\mathcal{K}}$, then it has finite index in $G_{\mathcal{K}}$.*

Proof. By reasonability $\text{Flim}(\mathcal{K}^*) = (\text{Flim}(\mathcal{K}), <^*)$ for some order $<^*$. Consider the logic action of $G_{\mathcal{K}}$ on LO (Example 2.4). Then $G_{\mathcal{K}^*}$ is the stabiliser of $<^*$. Hence, $|G_{\mathcal{K}} : G_{\mathcal{K}^*}| = |G_{\mathcal{K}} \cdot <^*|$ and it suffices to show that $G_{\mathcal{K}} \cdot <^*$ is finite. If $G_{\mathcal{K}^*}$ is normal, then it fixes every $R \in G_{\mathcal{K}} \cdot <^*$. Thus every such R is a union of 2-orbits. As $G_{\mathcal{K}^*}$ is oligomorphic, there are only finitely many such R . ■

We use the following mode of speech from [1]: Let \mathcal{K} be a Fraïssé class in the language L ; a *companion* of \mathcal{K} is a reasonable ordered Fraïssé class \mathcal{K}^* in the language $L \cup \{<\}$ which is Ramsey, has the ordering property and has L -reduct \mathcal{K} . Note:

PROPOSITION 4.10. *If a Fraïssé class is Ramsey, then it has a companion.*

Proof. By Lemmas 4.4 and 4.2. ■

PROPOSITION 4.11. *Let \mathcal{K} be a relational Fraïssé class that has a companion. If $M(G_{\mathcal{K}})$ is finite, then $|M(G_{\mathcal{K}})|$ is a power of 2.*

Proof. Let L denote the relational language of \mathcal{K} , and let \mathcal{K}^* be a companion of \mathcal{K} . By Theorem 2.8 we deduce that $F^* := \text{Flim}(\mathcal{K}^*) = (F, <^*)$ for $F := \text{Flim}(\mathcal{K})$, and that $M(G_{\mathcal{K}})$ is $\overline{G_{\mathcal{K}} \cdot <^*}$. Assume that $M(G_{\mathcal{K}})$ is finite. Then $G_{\mathcal{K}} \cdot <^*$ is finite, and since $G_{\mathcal{K}^*}$ is the stabiliser of $<^*$ in the logic action of $G_{\mathcal{K}}$ on LO , $G_{\mathcal{K}^*}$ has finite index in $G_{\mathcal{K}}$. By Theorem 2.6, $G_{\mathcal{K}^*}$ is extremely amenable. By Lemma 3.3, $G_{\mathcal{K}^*}$ is normal in $G_{\mathcal{K}}$ and $|M(G_{\mathcal{K}})| = |G_{\mathcal{K}} : G_{\mathcal{K}^*}|$.

Consider the diagonal actions of $G_{\mathcal{K}}$ and $G_{\mathcal{K}^*}$ on F^2 . We claim that for every $g \in G_{\mathcal{K}}$ and every 2-orbit S of $G_{\mathcal{K}^*}$ the set $g \cdot S \subseteq F^2$ is also a 2-orbit of $G_{\mathcal{K}^*}$. Indeed, normality implies that two pairs in the same 2-orbit of $G_{\mathcal{K}^*}$ are mapped by g to two pairs which are also in the same 2-orbit of $G_{\mathcal{K}^*}$, so there exists a 2-orbit T with $g \cdot S \subseteq T$. Reasoning analogously for g^{-1} and T we obtain $g^{-1} \cdot T \subseteq S$, so $g \cdot S = T$.

Call a 2-orbit S of $G_{\mathcal{K}^*}$ *black* if $a <^* b$ for all $(a, b) \in S$, and *white* if $b <^* a$ for all $(a, b) \in S$; orbits which are neither black nor white contain only pairs (a, b) with $a = b$. Let S be black or white. For every $g \in G_{\mathcal{K}}$, also $g(S)$ is black or white, and if $g(S)$ has the same colour as S , then $g(S) = S$. Indeed, as $g \in G_{\mathcal{K}}$, $g \upharpoonright \{a, b\}$ preserves all relations from L , and as $g(S)$ has the same colour as S , it also preserves $<^*$. Hence, for every $(a, b) \in S$, $g \upharpoonright \{a, b\}$ is a partial isomorphism of F^* , so it extends to some $h \in G_{\mathcal{K}^*}$ by homogeneity. Thus $g \cdot (a, b) = h \cdot (a, b)$, so $g \cdot (a, b) \in S$ and $g(S) = S$ follows.

We claim that $g^2 \in G_{\mathcal{K}^*}$ for every $g \in G_{\mathcal{K}}$. Seeking for contradiction, assume that there is an $(a, b) \in F^2$ such that $a <^* b$ is not equivalent to $g^2(a) <^* g^2(b)$. Then there is a black or white 2-orbit S of $G_{\mathcal{K}^*}$ such that $g^2(S)$ has a different colour. The colour of $g(S)$ equals that of S or $g^2(S)$, and consequently $S = g(S)$ or $g(S) = g^2(S)$. The first case $S = g(S)$ is impossible, because it implies $S = g^2(S)$. The second case $g(S) = g^2(S)$ is also impossible, because it implies the first via $g(S) = g^{-1}(g^2(S)) = g^{-1}(g(S)) = S$.

It follows that $G_{\mathcal{K}}/G_{\mathcal{K}^*}$ is an elementary abelian 2-group, i.e., it is the direct product of copies of the 2-element group. ■

THEOREM 4.12. *Let \mathcal{K} be a relational Fraïssé class with companion \mathcal{K}^* . Assume that $G_{\mathcal{K}}$ is oligomorphic. Then the following are equivalent:*

- (1) $|G_{\mathcal{K}} : G_{\mathcal{K}^*}|$ is finite.
- (2) $|G_{\mathcal{K}} : G_{\mathcal{K}^*}|$ is a finite power of 2.
- (3) $M(G_{\mathcal{K}})$ is finite.

- (4) $|M(G_{\mathcal{K}})|$ is a finite power of 2.
 (5) $G_{\mathcal{K}^*}$ is normal in $G_{\mathcal{K}}$.

Proof. By Theorem 2.8 we find that $F^* := \text{Flim}(\mathcal{K}^*) = (F, <^*)$ for $F := \text{Flim}(\mathcal{K})$, and that $M(G_{\mathcal{K}})$ is $\overline{G_{\mathcal{K}} \cdot <^*}$. Then $G_{\mathcal{K}^*}$ is oligomorphic by Lemma 4.8, and extremely amenable by Theorem 2.6. In a Hausdorff space a finite set equals its closure. As the elements of $G_{\mathcal{K}} \cdot <^*$ are in a one-to-one correspondence with $G_{\mathcal{K}}/G_{\mathcal{K}^*}$, we obtain (1) \Leftrightarrow (3) and (2) \Leftrightarrow (4). Proposition 4.11 implies (3) \Leftrightarrow (4), thus the first four items are equivalent. (5) \Rightarrow (1) follows from Lemma 4.9, and Lemma 3.3 implies (1) \Rightarrow (5). ■

COROLLARY 4.13. *Let \mathcal{K} be a relational Fraïssé class that has a companion. Then the Ramsey degree for embeddings of \mathcal{K} is either infinite or a finite power of 2.*

Proof. By Proposition 4.11 and Theorem 3.1. ■

Proof of Theorem 3.11. By Proposition 4.10 and Corollary 4.13. ■

Acknowledgements. We thank the anonymous referees for pointing out [25] and [15].

We thank Manuel Bodirsky, Lionel Nguyen Van Thé and Lyubomyr Zdomsky for their many comments on the manuscript.

The second author has received funding from the European Research Council under the European Community's Seventh Framework Programme (FP7/2007–2013 Grant Agreement no. 257039).

References

- [1] O. Angel, A. S. Kechris and R. Lyons, *Random orderings and unique ergodicity of automorphism groups*, J. Eur. Math. Soc. 16 (2014), 2059–2095.
- [2] H. Becker and A. S. Kechris, *The Descriptive Set Theory of Polish Group Actions*, London Math. Soc. Lecture Note Ser. 232, Cambridge Univ. Press, 1996.
- [3] M. Bodirsky, *New Ramsey classes from old*, Electron. J. Combin. 21 (2014), no. 2, Paper 2.22.
- [4] M. Bodirsky and M. Pinsker, *Reducts of Ramsey structures*, in: Model Theoretic Methods in Finite Combinatorics, Contemp. Math. 558, Amer. Math. Soc., 2011, 489–519.
- [5] M. Bodirsky, M. Pinsker and T. Tsankov, *Decidability of definability*, in: Proc. 26th Annual IEEE Sympos. on Logic in Computer Science (LICS'11), IEEE Computer Soc., 2011, 321–328.
- [6] P. J. Cameron, *Aspects of infinite permutation groups*, in: Groups St. Andrews 2005, Vol. 1, London Math. Soc. Lecture Note Ser. 339, Cambridge Univ. Press, 2007, 1–35.
- [7] G. Cherlin, *Two problems on homogeneous structures revisited*, in: Model Theoretic Methods in Finite Combinatorics, Contemp. Math. 558, Amer. Math. Soc., 2011, 319–415.

- [8] W. L. Fouché, *Symmetry and the Ramsey degree of posets*, Discrete Math. 167–168 (1997), 309–315.
- [9] W. L. Fouché, *Symmetries and Ramsey properties of trees*, Discrete Math. 197–198 (1999), 325–330.
- [10] W. L. Fouché, *Symmetry and the Ramsey degree of finite relational structures*, J. Combin. Theory Ser. A 85 (1999), 135–147.
- [11] R. L. Graham, B. L. Rothschild and J. H. Spencer, *Ramsey Theory*, Wiley-Intersci. Ser. Discrete Math. Optim., Wiley, 2nd ed., 1990.
- [12] M. Junker and M. Ziegler, *The 116 reducts of $(\mathbb{Q}, <, a)$* , J. Symbolic Logic 73 (2008), 861–884.
- [13] A. S. Kechris, *Dynamics of non-archimedean Polish groups*, in: Proc. 6th European Congress of Mathematics (Kraków, 2012), Eur. Math. Soc., 2014, 375–397.
- [14] A. S. Kechris, V. Pestov and S. Todorcevic, *Fraïssé limits, Ramsey theory, and topological dynamics of automorphism groups*, Geom. Funct. Anal. 15 (2005), 106–189.
- [15] A. S. Kechris and M. Sokić, *Dynamical properties of the automorphism groups of the random poset and random distributive lattice*, Fund. Math. 218 (2012), 69–94.
- [16] D. Macpherson, *A survey of homogeneous structures*, Discrete Math. 311 (2011), 1599–1634.
- [17] J. Nešetřil, *Ramsey theory*, in: Handbook of Combinatorics, Vol. 2, Elsevier, 1995, 1331–1403.
- [18] J. Nešetřil, *Ramsey classes and homogeneous structures*, Combin. Probab. Comput. 14 (2005), 171–189.
- [19] L. Nguyen Van Thé, *More on the Kechris–Pestov–Todorcevic correspondence: Precompact expansions*, Fund. Math. 222 (2013), 19–47.
- [20] L. Nguyen Van Thé, *Universal flows of closed subgroups of S_∞ and relative extreme amenability*, in: Asymptotic Geometric Analysis, Fields Inst. Comm. 68, Springer, 2013, 229–245.
- [21] V. Pestov, *Dynamics of Infinite-Dimensional Groups and Ramsey-Type Phenomena*, Publ. Mat. IMPA, Inst. Nac. Mat. Pura Apl., Rio de Janeiro, 2005.
- [22] K. Tent and M. Ziegler, *A Course in Model Theory*, Lecture Notes in Logic 40, Cambridge Univ. Press, 2012.
- [23] V. Uspenskij, *Compactifications of topological groups*, in: Proc. Ninth Prague Topological Sympos. (2001), Topology Atlas, North Bay, ON, 2002, 331–346.
- [24] A. Zucker, *Amenability and unique ergodicity of automorphism groups of Fraïssé structures*, Fund. Math. 226 (2014), 41–62.
- [25] A. Zucker, *Topological dynamics of closed subgroups of S_∞* , arXiv:1404.5057 [math.LO] (2014).

Moritz Müller
 Kurt Gödel Research Center (KGRC)
 1090 Wien, Austria
 E-mail: moritz.mueller@univie.ac.at

András Pongrácz
 Laboratoire d'Informatique (LIX)
 École Polytechnique
 91128 Palaiseau, France
 E-mail: andras.pongracz@lix.polytechnique.fr

*Received 25 April 2014;
 in revised form 28 July 2014*